

D-Branes on Cones and Gauge/String Dualities


Igor Klebanov

Department of Physics
Princeton University

Lectures at PiTP 2006
IAS, Princeton


QCD and String Theory

- At short distances, must smaller than 1 fermi, the quark-antiquark potential is approximately Coulombic, due to the Asymptotic Freedom.
- At large distances the potential should be linear (Wilson) due to formation of confining flux tubes.



Flux Tubes in QCD


- These objects may be approximately described by the Nambu strings
(animation from lattice work by D. Leinweber et al, Univ. of Adelaide)
- The tubes are widely used, for example, in jet hadronization algorithms (the Lund String Model) where they snap through quark-antiquark creation.



Large N Gauge Theories

- Connection of gauge theory with string theory is strengthened in 't Hooft's generalization from 3 colors (SU(3) gauge group) to N colors (SU(N) gauge group).
- Make N large, while keeping the 't Hooft coupling $\lambda = g_{\text{YM}}^2 N$ fixed.
- The probability of snapping a flux tube by quark-antiquark creation (meson decay) is $1/N$. The string coupling is $1/N$.
- Yet, the planar diagrams needed in the large N limit are very difficult to sum explicitly.

D-Branes vs. Geometry



- Dirichlet branes (Polchinski) led string theory back to gauge theory in the mid-90's.
- A stack of N Dirichlet 3-branes realizes $\mathcal{N}=4$ supersymmetric $SU(N)$ gauge theory in 4 dimensions. It also creates a curved background of 10-d theory of closed superstrings (artwork by E. Imeroni)

$$ds^2 = \left(1 + \frac{L^4}{r^4}\right)^{-1/2} \left(- (dx^0)^2 + (dx^i)^2\right) + \left(1 + \frac{L^4}{r^4}\right)^{1/2} (dr^2 + r^2 d\Omega_5^2)$$

which for small r approaches $AdS_5 \times S^5$

- Successful matching of graviton absorption by D3-branes, related to 2-point function of stress-energy tensor in the SYM theory, with a gravity calculation in the 3-brane metric (IK; Gubser, IK, Tseytlin) was a precursor of the AdS/CFT correspondence.

Conformal Invariance

- In the $\mathcal{N}=4$ SU(N) SYM theory there are 3 adjoint chiral superfields Z^i coupled to the $\mathcal{N}=1$ SU(N) SYM theory with superpotential $\text{Tr } Z^1 [Z^2, Z^3]$.
- The Asymptotic Freedom is canceled by the extra fields; the beta function is exactly zero! Hence, the theory is invariant under scale transformations $x^\mu \rightarrow \lambda x^\mu$. It is also invariant under space-time inversions.
- Such a theory is called a Conformal Field Theory (CFT).
- The $\mathcal{N}=4$ SU(N) SYM is also invariant under the SU(4) R-symmetry. Its full super-conformal symmetry is SU(2,2|4).

The AdS/CFT duality

Maldacena; Gubser, IK, Polyakov; Witten

- Relates conformal gauge theory in 4 dimensions to string theory on 5-d Anti-de Sitter space times a 5-d compact space. For the $\mathcal{N}=4$ SYM theory this compact space is a 5-sphere realizing the SU(4) R-symmetry.
- The SO(2,4) geometrical symmetry of the AdS₅ space realizes the conformal symmetry of the gauge theory.
- The d-dimensional AdS space is a hyperboloid

$$(X^0)^2 + (X^d)^2 - \sum_{i=1}^{d-1} (X^i)^2 = L^2 .$$

- Its metric is

$$ds^2 = \frac{L^2}{z^2} \left(dz^2 - (dx^0)^2 + \sum_{i=1}^{d-2} (dx^i)^2 \right)$$

- When a gauge theory is strongly coupled, the radius of curvature of the dual AdS_5 and of the 5-d compact space becomes large: $\frac{L^2}{\alpha'} \sim \sqrt{g_{\text{YM}}^2 N}$
- String theory in such a weakly curved background can be studied in the effective (super)-gravity approximation, which allows for a host of explicit calculations. Corrections to it proceed in powers of $\frac{\alpha'}{L^2} \sim \lambda^{-1/2}$
- Feynman graphs instead develop a weak coupling expansion in powers of λ . At weak coupling the dual string theory becomes difficult.

- Gauge invariant operators in the CFT_4 are in one-to-one correspondence with fields (or extended objects) in AdS_5
- Operator dimension is determined by the mass of the dual field; e.g. for scalar operators

$$\Delta_{\pm} = 2 \pm \sqrt{4 + m^2 L^2}$$

- Correlation functions are calculated from the dependence of string theory path integral on boundary conditions ϕ_0 in AdS_5 , imposed near $z=0$:


$$\langle \exp \int d^4 x \phi_0 \mathcal{O} \rangle = Z_{\text{string}}[\phi_0]$$

- In the large N limit the path integral is found from the classical string action: $Z_{\text{string}}[\phi_0] \sim \exp(-I[\phi_0])$

Conebrane Dualities

- To reduce the number of supersymmetries in AdS/CFT, we may place the stack of N D3-branes at the tip of a 6-d Ricci-flat cone X whose base is a 5-d Einstein space Y :

$$ds_X^2 = dr^2 + r^2 ds_Y^2$$



- Taking the near-horizon limit of the background created by the N D3-branes, we find the space $AdS_5 \times Y$, with N units of RR 5-form flux, whose radius is given by
- This type IIB background is conjectured to be dual to the IR limit of the gauge theory on N D3-branes at the tip of the cone X .

$$L^4 = \frac{\sqrt{\pi} \kappa N}{2 \text{Vol}(Y)} = 4\pi g_s N \alpha'^2 \frac{\pi^3}{\text{Vol}(Y)}$$

Trace Anomaly

- In a 4-d CFT there are two trace anomaly coefficients, a and c :

$$\langle T_{\alpha}^{\alpha} \rangle = -aE_4 - cI_4$$

$$E_4 = \frac{1}{16\pi^2} (R_{ijkl}^2 - 4R_{ij}^2 + R^2)$$
$$I_4 = -\frac{1}{16\pi^2} (R_{ijkl}^2 - 2R_{ij}^2 + \frac{1}{3}R^2)$$

- Calculations on $AdS_5 \times Y$ give their leading large N values

Henningson, Skenderis; Gubser

$$a = c = \frac{\pi^3 N^2}{4 \text{Vol}(Y)}$$

- In super-conformal theories the anomalies are related to the spectrum of R-charges of the chiral fermions:

Anselmi, Freedman, Grisaru, Johansen

$$a = \frac{3}{32} (3\text{Tr}R^3 - \text{Tr}R) ; \quad c = \frac{1}{32} (9\text{Tr}R^3 - 5\text{Tr}R)$$

- This provides basic checks of the dualities.
- For the $\mathcal{N}=4$ SYM theory the gauginos have $R=1$, while the fermion fields from the Z^i chiral multiplets have $R=-1/3$.
- Since $\text{Tr } R=0$, we find $a=c$, and

$$a = c = \frac{9}{32} \text{Tr} R^3 = (N^2 - 1) \frac{9}{32} (1 + 3(-1/3)^3) = \frac{N^2 - 1}{4}$$

- On the gravity side, the volume of S^5 is π^3 (the radius of Y is fixed so that $R_{ij} = 4g_{ij}$)
- For large N the two calculations of anomaly coefficients agree.

Orbifold Cones

Kachru, Silverstein; Lawrence, Nekrasov, Vafa

- The simplest set of examples is provided by cones that are orbifolds R^6/Γ , where Γ is a subgroup of the rotation group $SU(4)$.
- For abelian orbifolds, all group elements can be brought to the form
- For Z_k orbifolds, the n -th group element is specified by three integers m_i defined mod k : $x_i = nm_i/k$.
- If none of the eigenvalues of the generator = 1, then all SUSY is broken; if one of the eigenvalues = 1, then $\mathcal{N}=1$ SUSY is preserved; if two of the eigenvalues = 1, then $\mathcal{N}=2$ SUSY is preserved.

$$\begin{pmatrix} e^{2\pi i x_1} & 0 & 0 & 0 \\ 0 & e^{2\pi i x_2} & 0 & 0 \\ 0 & 0 & e^{2\pi i x_3} & 0 \\ 0 & 0 & 0 & e^{-2\pi i(x_1+x_2+x_3)} \end{pmatrix}$$


- The action in the n -th twisted sector on 3 complex coordinates of C^3 , $Z^1 = X^1 + iX^2$, $Z^2 = X^3 + iX^4$, $Z^3 = X^5 + iX^6$

and their complex conjugates, is

$$R(g_n) = \text{diag}(\omega_k^{n(m_1+m_2)}, \omega_k^{n(m_1+m_3)}, \omega_k^{n(m_2+m_3)}, \omega_k^{-n(m_1+m_2)}, \omega_k^{-n(m_1+m_3)}, \omega_k^{-n(m_2+m_3)})$$

where $\omega_k = e^{2\pi i/k}$

- If none of these phases = 1, then the orbifold acts freely on S^5/Γ . (The tip of the cone is a fixed point that is removed in the basic near-horizon limit.)
- A well-known example of a freely-acting orbifold is Z_3 with $m_i=1$. Since one of the eigenvalues of the generator = 1, i.e. $\Gamma \subset SU(3)$, this orbifold preserves $\mathcal{N}=1$ SUSY.



Construction of the quiver gauge theories

Douglas, Moore

- Gauge theory on N D3-branes at the tip of R^6/Γ is found by applying projections to the $U(Nk)$ gauge theory on the covering space. Retain only the fields invariant under the orbifold action combined with conjugation by a $U(Nk)$ matrix γ acting on the gauge indices:

$$\gamma = \text{diag}(I_N, e^{2\pi i/k} I_N, e^{4\pi i/k} I_N, \dots, e^{-2\pi i/k} I_N)$$

$$\psi^1 \rightarrow e^{2\pi i m_1/k} \gamma \psi^1 \gamma^{-1}, \quad \psi^2 \rightarrow e^{2\pi i m_2/k} \gamma \psi^2 \gamma^{-1}, \dots$$

$$Z^1 \rightarrow e^{2\pi i(m_1+m_2)/k} \gamma Z^1 \gamma^{-1}, \quad Z^2 \rightarrow e^{2\pi i(m_1+m_3)/k} \gamma Z^2 \gamma^{-1}, \dots$$

- In the supersymmetric examples, such as C^3/Z_3 or the conifold, the conebrane dualities have been tested almost as thoroughly as in the maximally supersymmetric case.
- But when all SUSY is broken, problems may arise.
- When the orbifold Γ breaks all SUSY and is not freely acting, then the weakly curved background $AdS_5 \times S^5/\Gamma$ is unstable due to the presence of tachyons that have $(mL)^2 < -4$, and therefore violate the BF bound.
- But for freely acting orbifolds, the negative zero-point energy is compensated by the large stretching of the twisted sector closed strings in the compact space. Hence, at large radius, there are no 'bad tachyons.' This makes freely acting orbifolds particularly interesting from AdS/CFT point of view.
- Yet, before the formal decoupling limit, the non-SUSY freely acting orbifolds have closed-string tachyons localized at the tip of the cone.

Closed String Zero-Point Energy

- In light-Cone Green-Schwarz, there are 4 complex world sheet bosons and fermions.

- In the n-th twisted sector the boundary conditions on the

fermions are $b^l(\sigma + 2\pi) = e^{2\pi i n m_l / k} b^l(\sigma)$

and on the bosons

$$X^1(\sigma + 2\pi) = X^1(\sigma) ,$$

$$X^2(\sigma + 2\pi) = e^{2\pi i n (m_1 + m_2) / k} X^2(\sigma)$$

$$X^3(\sigma + 2\pi) = e^{2\pi i n (m_1 + m_3) / k} X^3(\sigma)$$

$$X^4(\sigma + 2\pi) = e^{2\pi i n (m_2 + m_3) / k} X^4(\sigma)$$

- The total zero-point energy of these modes is

$$8E_0(x_1, x_2, x_3) = -1 - (2\{x_1 + x_2\} - 1)^2 - (2\{x_1 + x_3\} - 1)^2 - (2\{x_2 + x_3\} - 1)^2 \\ + (2\{x_1 + x_2 + x_3\} - 1)^2 + \sum_{i=1}^3 (2\{x_i\} - 1)^2$$

Weak coupling analysis of non-SUSY quivers

- My recent work with Dymarsky and Roiban (hep-th/0505099 and 0509132) reconsiders quiver gauge theory on a stack of D3-branes at the tip of a cone R^6/Γ where the orbifold group Γ breaks all the supersymmetry.
- At first sight, the gauge theory seems conformal because the planar beta functions for all single-trace operators vanish. The candidate string dual is $AdS_5 \times S^5/\Gamma$. Kachru, Silverstein; Lawrence, Nekrasov, Vafa; Bershadsky, Johanson
- However, dimension 4 double-trace operators made out of twisted single-trace ones, $f O_n O_{-n}$, are induced at one-loop. Their planar beta-functions have the form

$$\beta_f = a \lambda^2 + 2 \gamma f \lambda + f^2$$

$$\beta_\lambda = 0$$

A Note on Normalizations

- The VEV of a single trace operator is of order N .
- The standard Yang-Mills action $S = - \int d^4x \frac{1}{2g_{YM}^2} \text{Tr} F_{\mu\nu}^2$ is of order N^2 in the 't Hooft limit.
- The double-trace operators $f O_n O_{-n}$ make contributions of the same order (for the coupling constant of order 1). They cannot be ignored in the leading large N limit.
- In fact, the tree-level potential of the $SU(N)^k$ quiver theories (with the interacting $U(1)$'s decoupled) contains such double-trace terms.

Problem 1


a) Derive the Z_2 quiver gauge theory obtained by the projection on the $U(2N)$ $\mathcal{N} = 4$ SYM where the Z_2 is generated by $-I$ in $SU(4)$ accompanied by conjugation with $\gamma = \text{diag}(I_N, -I_N)$. What is the gauge group and field content? Is this a freely acting orbifold?

b) In this gauge theory, calculate the one-loop Coleman-Weinberg potential as a function of the eigenvalues of the adjoint scalars. What are the operators that pick up one-loop beta functions?


- If $D = \gamma^2 - a < 0$, then there is no real fixed point for f .
- A class of Z_k orbifolds with global $SU(3)$ symmetry, that are freely acting on the 5-sphere, has the group action in the fundamental of $SU(4)$

$$r(g^n) = \text{diag}(\omega_k^n, \omega_k^n, \omega_k^n, \omega_k^{-3n})$$

$$\omega_k = e^{i\alpha_k}, \quad \alpha_k = \frac{2\pi}{k}$$



- Here is a plot of a one-loop $SU(N)^k$ gauge theory discriminant, D , and of the ground state closed string m^2 on the cone without the D-branes. $n=1, \dots, k-1$ labels the twisted sector, and $x=n/k$.
- The simplest freely acting non-susy example is Z_5 where there are four induced double-trace couplings



$$\delta_2 \text{ trace } \mathcal{L} = f_{8,1} O_1^{(i\bar{j})} O_{-1}^{(j\bar{i})} + f_{8,2} O_2^{(i\bar{j})} O_{-2}^{(j\bar{i})} + f_{1,1} O_1 O_{-1} + f_{1,2} O_2 O_{-2}$$


- For example, the $SU(3)$ adjoints

$$O_n^{(i\bar{j})} \text{ are}$$


$$\sum_{k=1}^5 \left(\Phi_{k,k+2}^i \Phi_{k+2,k}^{\bar{j}} - \frac{1}{3} \eta^{i\bar{j}} \Phi_{k,k+2}^l \Phi_{k+2,k}^{\bar{l}} \right) e^{in\alpha(k-1)}$$

$$(\alpha = 2\pi/5)$$

- For more complicated orbifolds, crossing of eigenvalues of the discriminant matrix becomes important. The agreement with closed strings continues to hold.



- Generally, there are three twist angles x_i that define a cube. The stability/instability regions agree between one-loop gauge theory and string theory.



- Any non-SUSY abelian orbifold contains unstable operators. This appears to remove all such orbifold quivers from a list of large N perturbatively conformal gauge theories.
- The one-loop beta functions destroy the conformal invariance precisely in those twisted sectors where there exist closed-string tachyons localized at the tip of R^6/Γ . Thus, a very simple correspondence emerges between perturbative gauge theory and free closed string on an orbifold. Why? Perhaps, in the presence of tachyons, the standard AdS/CFT decoupling argument may fail.
- The $AdS_5 \times S^5/\Gamma$ background is tachyon-free at large radius. Could it have some instabilities? If not, then there is a transition from instability to stability as λ is increased.

- What is the end-point of the RG flow?
- Condensation of localized tachyon smoothes out the tip of the cone. Adams, Polchinski, Silverstein
- The gauge theory on D3-branes at a smooth point is $\mathcal{N}=4$ SYM. Hence, a natural conjecture is that the gauge theory flows from the non-SUSY $SU(N)^k$ quiver gauge theory to the $\mathcal{N}=4$ $SU(N)$ SYM. Dymarsky, Franco, Roiban, IK (work in progress)

D3-branes on the Conifold

- The conifold is a Calabi-Yau 3-fold cone X described by the constraint $\sum_{a=1}^4 z_a^2 = 0$ on 4 complex variables.
- Its base Y is a coset $T^{1,1}$ which has symmetry $SU(2)_A \times SU(2)_B$ that rotates the z 's, and also $U(1)_R$: $z_a \rightarrow e^{i\theta} z_a$

- The Sasaki-Einstein metric on $T^{1,1}$ is

$$ds_{T^{1,1}}^2 = \frac{1}{9} \left(d\psi + \sum_{i=1}^2 \cos \theta_i d\phi_i \right)^2 + \frac{1}{6} \sum_{i=1}^2 \left(d\theta_i^2 + \sin^2 \theta_i d\phi_i^2 \right)$$

where $\theta_i \in [0, \pi], \phi_i \in [0, 2\pi], \psi \in [0, 4\pi]$

- The topology of $T^{1,1}$ is $S^2 \times S^3$.

- The $\mathcal{N}=1$ SCFT on the D3-branes at the apex of the conifold has gauge group $SU(N) \times SU(N)$ coupled to bifundamental chiral superfields A_1, A_2 , in $(\bar{\mathbf{N}}, \mathbf{N})$, and B_1, B_2 in $(\mathbf{N}, \bar{\mathbf{N}})$. IK, Witten
- The R-charge of each field is $1/2$. This insures $U(1)_R$ anomaly cancellation.
- The unique $SU(2)_A \times SU(2)_B$ invariant, exactly marginal quartic superpotential is added:

$$W = \epsilon^{ij} \epsilon^{kl} \text{tr} A_i B_k A_j B_l$$

- This theory also has a baryonic $U(1)$ symmetry under which $A_k \rightarrow e^{ia} A_k$; $B_l \rightarrow e^{-ia} B_l$, and a Z_2 symmetry which interchanges the A's with the B's and implements charge conjugation.

Comparison with a Z_2 Orbifold Quiver

- The simplest $\mathcal{N}=2$ SUSY quiver has $k=2$; $m_1=m_2=1$, $m_3=0$. The gauge group is again $SU(N) \times SU(N)$, but in addition to the bifundamentals A_i , B_j , there is one adjoint chiral superfield for each gauge group, with superpotential $g\text{Tr}\Phi(A_1B_1 - A_2B_2) + g\text{Tr}\tilde{\Phi}(B_1A_1 - B_2A_2)$

- Adding a Z_2 odd mass term $\frac{m}{2}(\text{Tr}\Phi^2 - \text{Tr}\tilde{\Phi}^2)$ and integrating out the adjoints, we obtain the superpotential of the conifold theory,

$$-\frac{g^2}{m} [\text{Tr}(A_1B_1A_2B_2) - \text{Tr}(B_1A_1B_2A_2)]$$

Problem 2

In a supersymmetric field theory, the trace anomaly coefficients a and c are given by the formulae

$$a = \frac{3}{32} (3\text{Tr}R^3 - 3\text{Tr}R) , \quad c = \frac{1}{32} (9\text{Tr}R^3 - 5\text{Tr}R) ,$$

where R refers to the $U(1)_R$ charges, and the trace is over all the chiral fermion fields.

a) Calculate a and c in the following two gauge theories: the $\mathcal{N} = 2$ supersymmetric Z_2 orbifold quiver, and in the $\mathcal{N} = 1$ SCFT on N D3-branes at the conifold.

b) For $AdS_5 \times Y$ with N units of RR 5-form flux, it was found at leading order in N that

$$a = c = \frac{N^2 \pi^3}{4\text{vol}(Y)} ,$$

where the radius of Y is normalized so that $R_{ij} = 4g_{ij}$ on Y . Compare this formula with the gauge theory results of part a).


Breaking the Conformal Symmetry

- A useful tool is to add to the N D3-branes M D5-branes wrapped over the S^2 at the tip of the conifold.
- The 10-d geometry dual to the gauge theory on these branes is the warped deformed conifold (IK, Strassler)

$$ds_{10}^2 = h^{-1/2}(t)(-(dx^0)^2 + (dx^i)^2) + h^{1/2}(t)ds_6^2$$

- ds_6^2 is the metric of the deformed conifold, a simple Calabi-Yau space defined by the following constraint on 4 complex variables:

$$\sum_{i=1}^4 z_i^2 = \epsilon^2$$




String Theoretic Approach to Confinement

- It is possible to generalize the AdS/CFT correspondence in such a way that the quark-antiquark potential is linear at large distance.
- A "cartoon" of the necessary metric is

$$ds^2 = \frac{dz^2}{z^2} + a^2(z)(-(dx^0)^2 + (dx^i)^2)$$

- The space ends at a maximum value of z where the warp factor is finite. Then the confining string tension is

$$\frac{a^2(z_{\max})}{2\pi\alpha'}$$



- The warp factor is finite at the `end of space' $t=0$, as required for the confinement: $h(t) = 2^{-8/3} \gamma I(t)$

$$I(t) = \int_t^\infty dx \frac{x \coth x - 1}{\sinh^2 x} (\sinh 2x - 2x)^{1/3}, \quad \gamma = 2^{10/3} (g_s M \alpha')^2 \varepsilon^{-8/3}$$

- The standard warp factor a^2 , which measures the string tension, is identified with $h(t)^{-1/2}$ and is minimized at $t=0$. It blows up at large t (near the boundary).
- The dilaton is exactly constant due to the self-duality of the 3-form background

$$\star_6 G_3 = iG_3, \quad G_3 = F_3 - \frac{i}{g_s} H_3$$

- The radius-squared of the S^3 at $t=0$ is $g_s M$ in string units.
- When $g_s M$ is large, the curvatures are small everywhere, and the SUGRA solution is reliable in 'solving' this confining gauge theory.

Anomalously light glueballs

- The confining string tension is $T_s = \frac{1}{2^{4/3} a_0^{1/2} \pi} \frac{\epsilon^{4/3}}{(\alpha')^2 g_s M}$
- The glueballs are the normalizable modes localized near at small t . In the supergravity limit (at large $g_s M$) their mass scales are

$$m_{glueball} \sim m_{KK} \sim \frac{\epsilon^{2/3}}{g_s M \alpha'}$$

$$T_s \sim g_s M (m_{glueball})^2$$

- In order to eliminate the anomalously light bound states, we need a small $g_s M$, which requires a departure from the SUGRA limit.
- Even for small $g_s M$, SUGRA becomes reliable in the UV (at large t).

Log running of couplings

- The large radius asymptotic solution is characterized by logarithmic deviations from $AdS_5 \times T^{1,1}$ IK, Tseytlin

- The near-AdS radial coordinate is $r \sim \varepsilon^{2/3} e^{t/3}$

- The NS-NS and R-R 2-form potentials:

$$F_3 = \frac{M\alpha'}{2}\omega_3, \quad B_2 = \frac{3g_s M\alpha'}{2}\omega_2 \ln(r/r_0)$$

$$\omega_2 = \frac{1}{2}(g^1 \wedge g^2 + g^3 \wedge g^4) = \frac{1}{2}(\sin \theta_1 d\theta_1 \wedge d\phi_1 - \sin \theta_2 d\theta_2 \wedge d\phi_2)$$

$$\omega_3 = \frac{1}{2}g^5 \wedge (g^1 \wedge g^2 + g^3 \wedge g^4)$$

- This translates into log running of the gauge couplings through
- The warp factor deviates from the $M=0$ solution logarithmically.

$$\frac{4\pi^2}{g_1^2} + \frac{4\pi^2}{g_2^2} = \frac{\pi}{g_s e^\Phi},$$

$$\left[\frac{4\pi^2}{g_1^2} - \frac{4\pi^2}{g_2^2} \right] g_s e^\Phi = \frac{1}{2\pi\alpha'} \left(\int_{S^2} B_2 \right) - \pi$$

$$h(r) = \frac{27\pi(\alpha')^2 [g_s N + a(g_s M)^2 \ln(r/r_0) + a(g_s M)^2/4]}{4r^4}$$


- Remarkably, the 5-form flux, dual to the number of colors, also changes logarithmically with the RG scale.

$$\tilde{F}_5 = \mathcal{F}_5 + \star \mathcal{F}_5, \quad \mathcal{F}_5 = 27\pi\alpha'^2 N_{eff}(r) \text{vol}(T^{1,1})$$

$$N_{eff}(r) = N + \frac{3}{2\pi} g_s M^2 \ln(r/r_0)$$

- What is the explanation in the dual $SU(kM) \times SU((k-1)M)$ SYM theory coupled to bifundamental chiral superfields A_1, A_2, B_1, B_2 ? A novel phenomenon, called a **duality cascade**, takes place: k repeatedly changes by 1 as a result of the Seiberg duality IK, Strassler

(diagram of RG flows from a review by M. Strassler)



$Y^{p,q}$ Dualities

- Cascading behavior is not limited to the conifold. Recently, an infinite family of new CY cones over Sasaki-Einstein spaces $Y^{p,q}$ of topology $S^2 \times S^3$ have been constructed (p and q are coprime integers). Gauntlett, Martelli, Sparks, Waldram

$$d\Omega_{Y^{p,q}}^2 = (e^\theta)^2 + (e^\phi)^2 + (e^y)^2 + (e^\beta)^2 + (e^\psi)^2$$

$$e^\theta = \sqrt{\frac{1-y}{6}} d\theta, \quad e^\phi = \sqrt{\frac{1-y}{6}} \sin\theta d\phi$$

$$e^y = \frac{1}{\sqrt{wv}} dy, \quad e^\beta = \frac{\sqrt{wv}}{6} (d\beta + \cos\theta d\phi)$$

$$e^\psi = \frac{1}{3} (d\psi - \cos\theta d\phi + y(d\beta + \cos\theta d\phi))$$

$$w(y) = \frac{2(b-y^2)}{1-y},$$

$$v(y) = \frac{b-3y^2+2y^3}{b-y^2}$$

$$b = \frac{1}{2} - \frac{p^2 - 3q^2}{4p^3} \sqrt{4p^2 - 3q^2}$$

- y ranges between two smaller roots of $v(y)$:

$$y_1 = \frac{1}{4p} \left(2p - 3q - \sqrt{4p^2 - 3q^2} \right)$$


$$y_2 = \frac{1}{4p} \left(2p + 3q - \sqrt{4p^2 - 3q^2} \right)$$

$$\text{Vol}(Y^{p,q}) = \frac{q^2 \left(2p + \sqrt{4p^2 - 3q^2} \right)}{3p^2 \left(3q^2 - 2p^2 + p\sqrt{4p^2 - 3q^2} \right)} \pi^3$$

- The $SU(N)^{2p}$ SCFT's on N D3-branes at the tip of the cones have also been constructed.

Benvenuti, Franco, Hanany, Martelli, Sparks

- For example, here is the quiver diagram for the SCFT dual to $AdS_5 \times Y^{4,3}$



R-charges from a-maximization


- The conformal invariance conditions do not fully determine the R-charges. Let $R_Z=x$, $R_Y=y$, $R_U=1-(x+y)/2$, $R_V=1+(x-y)/2$
- The technique of a-maximization Intriligator, Wecht gives

$$\begin{aligned}x &= \frac{1}{3q^2} \left(-4p^2 + 2pq + 3q^2 + (2p - q) \sqrt{4p^2 - 3q^2} \right) \\y &= \frac{1}{3q^2} \left(-4p^2 - 2pq + 3q^2 + (2p + q) \sqrt{4p^2 - 3q^2} \right)\end{aligned}$$

- Remarkably, this gives the trace anomaly agreeing with the AdS/CFT

$$a(Y^{p,q}) = \frac{\pi^3 N^2}{4 \text{Vol}(Y^{p,q})}$$

- Addition of M D5-branes wrapped over the S^2 modifies the quiver diagram to
- Performing the Seiberg duality on the biggest gauge group gives the same quiver with $N \rightarrow N-M$. This fact is necessary for the existence of a self-similar cascading RG flow.
- The gravity duals of these cascades include the ISD 3-form field strength. Herzog, Ejaz, IK



$$\Omega_{2,1} = K \left(\frac{dr}{r} + ie^\psi \right) \wedge \omega$$

$$\omega = F(y)(e^\theta \wedge e^\phi - e^y \wedge e^\beta)$$

$$F(y) = \frac{1}{(1-y)^2}$$

- The metric and 5-form are determined by a single warp factor, which however depends on 2 coordinates.
- Luckily, the PDE for $h(r,y)$ is exactly solvable

$$ds^2 = h^{-1/2} dx_4^2 + h^{1/2} (dr^2 + r^2 d\Omega_{Y^{p,q}}^2)$$

$$g_s F_5 = d(h^{-1}) \wedge d^4 x + *[d(h^{-1}) \wedge d^4 x]$$

$$h(r, y) = \frac{A \ln(r/r_0) + s(y)}{r^4}$$

$$s(y) = -\frac{C}{4(b-1)} \left[\frac{1}{1-y} + \frac{(1+2y_1)(1+2y_2) \ln(y_3 - y)}{2(b-1)} \right]$$

- The log behavior characteristic of the cascade produces a naked singularity in the IR. Is there a smooth solution with the above large r behavior?

IR Behavior of the Conifold Cascade

- Here the dynamical deformation of the conifold renders the solution smooth, and explains the IR dynamics of the gauge theory.
- **Dimensional transmutation** in the IR. The dynamically generated confinement scale is

$$\sim \varepsilon^{2/3}$$

- The pattern of **R-symmetry breaking** is the same as in the SU(M) SYM theory: $Z_{2M} \rightarrow Z_2$
- Yet, the IR gauge theory is somewhat more complicated.

- In the IR the gauge theory cascades down to $SU(2M) \times SU(M)$. The $SU(2M)$ gauge group effectively has $N_f = N_c$.
- The baryon and anti-baryon operators Seiberg

$$\mathcal{A} = \epsilon^{i_1 \dots i_{N_c}} A_{\alpha_1 i_1}^{a_1} \dots A_{\alpha_{N_c} i_{N_c}}^{a_{N_c}}$$

$$\mathcal{B} = \epsilon_{i_1 \dots i_{N_c}} B_{\dot{\alpha}_1 a_1}^{i_1} \dots B_{\dot{\alpha}_{N_c} a_{N_c}}^{i_{N_c}}$$

acquire expectation values and break the $U(1)$ symmetry under which $A_k \rightarrow e^{ia} A_k$; $B_l \rightarrow e^{-ia} B_l$. Hence, we observe confinement without a mass gap: due to $U(1)_{\text{baryon}}$ **chiral symmetry breaking** there exist a Goldstone boson and its massless scalar superpartner.

- The KS solution is part of a moduli space of confining SUGRA backgrounds, **resolved warped deformed conifolds**. Gubser, Herzog, IK; Butti, Grana, Minasian, Petrini, Zaffaroni

- To look for them we need to use the PT ansatz:

$$ds_{10}^2 = H^{-1/2} dx_m dx_m + e^x ds_6^2,$$

$$ds_6^2 = (e^g + a^2 e^{-g})(e_1^2 + e_2^2) + e^{-g} \sum_{i=1}^2 (\epsilon_i^2 - 2a e_i \epsilon_i) + v^{-1}(\tilde{\epsilon}_3^2 + dt^2)$$

- H , x , g , a , v , and the dilaton are functions of the radial variable t .
- Additional radial functions enter into the ansatz for the 3-form field strengths. The PT ansatz preserves the $SO(4)$ but breaks a Z_2 charge conjugation symmetry, except at the KS point.

- BGMPZ used the method of SU(3) structures to derive the complete set of coupled first-order equations.
- A result of their integration is that the warp factor and the dilaton are related:

$$H(t) = \tilde{H} \left(e^{-2\phi(t)} - 1 \right)$$

Dymarsky, IK, Seiberg

- The integration constant determines the 'modulus' U: $\tilde{H} = \gamma U^{-2}$ where $\gamma = 2^{10/3} (g_s M \alpha')^2 \varepsilon^{-8/3}$
- At large t the solution approaches the KT 'cascade asymptotics':

$$a(t) = -2e^{-t} + Ue^{-5t/3}(-t + 1) + \dots$$

$$\gamma^{-1} H(t) = \frac{3}{32} e^{-4t/3} (4t - 1) - \frac{3}{32 \cdot 512} U^2 (256t^3 - 864t^2 + 1752t - 847) e^{-8t/3} + O\left(e^{-10t/3}\right)$$



- The resolution parameter U is proportional to the VEV of the operator

$$U = \text{Tr} \left(\sum_{\alpha} A_{\alpha} A_{\alpha}^{\dagger} - \sum_{\dot{\alpha}} B_{\dot{\alpha}}^{\dagger} B_{\dot{\alpha}} \right)$$

- This family of resolved warped deformed conifolds is dual to the 'baryonic branch' in the gauge theory (the quantum deformed moduli space):


$$\mathcal{A} = i\Lambda_1^{2M} \zeta, \quad \mathcal{B} = i\Lambda_1^{2M} / \zeta$$

- Various quantities have been calculated as a function of the modulus $U = \ln |\zeta|$.
 - Here are plots of the string tension (a **fundamental** string at the bottom of the throat is **dual to an 'emergent'** chromo-electric flux tube) and of the dilaton profiles
- Dymarsky, IK, Seiberg



BPS Domain Walls



- A D5-brane wrapped over the 3-sphere at the bottom of the throat is the domain wall separating two adjacent vacua of the theory.
- Since it is BPS saturated, its tension cannot depend on the baryonic branch modulus. This is indeed the case. This fact provides a check on the choice of the UV boundary conditions, and on the numerical integration procedure necessary away from the KS point.
- Analytic proof?



Applications to D-brane Inflation

- The Slow-Roll Inflationary Universe (Linde; Albrecht, Steinhardt) is a very promising idea for generating the CMB anisotropy spectrum observed by the WMAP.
- Finding models with very flat potentials has proven to be difficult. Recent string theory constructions use moving D-branes. Dvali, Tye, ...
- In the KKLT/KKLMMT model, the warped deformed conifold is embedded into a string compactification. An anti-D3-brane is added at the bottom to break SUSY and generate a potential. A D3-brane rolls in the throat. Its radial coordinate plays the role of an inflaton.


Kachru, Kallosh, Linde, Maldacena, McAllister, Trivedi



A related suggestion for D-brane inflation

(A. Dymarsky, IK, N. Seiberg)

- In a flux compactification, the $U(1)_{\text{baryon}}$ is gauged. Turn on a Fayet-Iliopoulos parameter ξ .
- This makes the throat a **resolved** warped deformed conifold.
- The probe D3-brane potential on this space is asymptotically flat, if we ignore effects of compactification and D7-branes. The plots are for two different values of $U \sim \xi$.
- No anti-D3 needed: in presence of the D3-brane, SUSY is broken by the D-term ξ . Related to the 'D-term Inflation' Binetruy, Dvali; Halyo




Slow roll D-brane inflation?

- Effects of D7-branes and of compactification generically spoil the flatness of the potential. Non-perturbative effects introduce the KKLT-type superpotential

$$W = W_0 + A(X)e^{-a\rho}$$

where X denotes the D3-brane position. In any warped throat D-brane inflation model, it is important to calculate $A(X)$.



- The gauge theory on D7-branes wrapping a 4-cycle Σ_4 has coupling $\frac{1}{g^2} = \frac{V_{\Sigma_4}^w}{g_7^2} = \frac{T_3 V_{\Sigma_4}^w}{8\pi^2}$

- The non-perturbative superpotential $\propto \exp\left(-\frac{T_3 V_{\Sigma_4}^w}{N_{D7}}\right)$

depends on the D3-brane location through the warped volume $V_{\Sigma_4}^w \equiv \int_{\Sigma_4} d^4\xi \sqrt{g^{ind}} h(X)$

- In the throat approximation, the warp factor can be calculated and integrated over a 4-cycle explicitly. Baumann, Dymarsky, Klebanov, Maldacena, McAllister, Murugan.

- For a class of conifold embeddings

Arean, Crooks, Ramallo ($w_1 = z_1 + iz_2$, etc.)

$$\prod_{i=1}^4 w_i^{p_i} = \mu^P$$

$$P \equiv \sum_{i=1}^4 p_i$$

the result is

$$A = A_0 \left(\frac{\mu^P - \prod_{i=1}^4 w_i^{p_i}}{\mu^P} \right)^{1/n}$$

- This formula applies both to n wrapped D7-branes, and to a wrapped Euclidean D3 ($n=1$).
- For the latter case, Ganor showed that A has a simple zero when the D3-brane approaches the 4-cycle. Our result agrees with this.
- We have also carried out such calculations for 4-cycles within the Calabi-Yau cones over $Y^{p,q}$ with analogous results: $A(X)$ is proportional to the embedding equation raised to the power $1/n$. This appears to be a general rule for 4-cycles in the throat.

- The dependence of the non-perturbative superpotential on D3-brane position, and other compactification effects, give Hubble-scale corrections to the inflaton potential.
- Some ‘fine-tuning’ is generally needed to cancel different corrections to the D3-brane potential. This is currently under investigation with D. Baumann, A. Dymarsky, J. Maldacena, L. McAllister and P. Steinhardt.

Conclusions

- In the first part, we investigated non-SUSY orbifolds of AdS/CFT. At one loop, flow of double-trace couplings spoils conformal invariance even in the large N limit. There is a precise connection of this instability with presence of twisted sector closed string tachyons.
- Gauge/string dualities for confining gauge theories give a new geometrical view of such important phenomena as dimensional transmutation, chiral symmetry breaking, and quantum deformation of moduli space.
- Embedding gauge/string dualities into string compactifications offers new possibilities for physics beyond the SM, and for modeling inflation. In particular, D3-branes on resolved warped deformed conifolds may realize D-term inflation.
- Calculation of non-perturbative corrections to the inflaton potential is important for determining if these models can produce slow-roll inflation.